

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/279761143>

Artocarpus heterophyllus (Jackfruit): An overview

Article in *Pharmacognosy Reviews* · July 2009

CITATIONS

173

READS

24,608

4 authors, including:



Om Prakash

Goel Institute of Pharmacy & Sciences, Lucknow

41 PUBLICATIONS 683 CITATIONS

[SEE PROFILE](#)



Anurag Mishra

Ashoka institute of technology & management

38 PUBLICATIONS 1,906 CITATIONS

[SEE PROFILE](#)



Rajiv Gupta

Babu Banarasi Das University

67 PUBLICATIONS 2,838 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Increasing solubility of poorly soluble drugs and anticancer activity of sesamol [View project](#)



For Antidiabetic Activity [View project](#)

PHCOG REV. : Review Article

Artocarpus heterophyllus (Jackfruit): An Overview

Om Prakash, Rajesh Kumar, Anurag Mishra, Rajiv Gupta*

Department of Pharmacognosy, Faculty of Pharmacy, Babu Banarasi Das National Institute of Technology & Management, Dr. Akhilesh Das Nagar, Lucknow 227 105, U.P., India

*Address for Correspondence: bbdcognosy@rediffmail.com, opverma2007@gmail.com

Mobile No. 09839278227

ABSTRACT

Artocarpus heterophyllus (Syn. Kathal) belonging to family Moraceae is an integral part of common Indian diet and is freely available in Indian and adjoining continents, its medicinal properties are also mentioned in Ayurveda. The plant is reported to possess antibacterial, anti-inflammatory, antidiabetic, antioxidant and immunomodulatory properties. *Artocarpus heterophyllus* is an important source of compounds like morin, dihydromorin, cynomacurin, artocarpin, isoartocarpin, cyloartocarpin, artocarpesin, oxydihydroartocarpesin, artocarpetin, norartocarpetin, cycloartinone, betulinic acid, artocarpanone and heterophyllol which are useful in fever, boils, wounds, skin diseases, convulsions, diuretic, constipation, ophthalmic disorders and snake bite etc.

KEY WORDS: *Artocarpus heterophyllus*, Moraceae, Phytochemistry, Pharmacological activities.

INTRODUCTION

The *Artocarpus heterophyllus* is a species of tree of the mulberry family (Moraceae) is known by other names jackfruit (Eng.), Kathal, Panas (Hindi), Kanthal (Beng.), Palaa (Tamil), Phanas (Guj & Mar) & Chakka (Malayalam). It is native to Western Ghats of India, Malaysia and also found in central and eastern Africa, south-eastern Asia, the Caribbean, Florida, Brazil, Australia, Puerto Rico and many Pacific Islands (1). It is a large, evergreen tree, 10-15m in height, indigenous to the evergreen forests at altitude of 450-1,200m and cultivated throughout the hotter parts of India. Stem of this plant is straight rough whereas bark is green or black, 1.25cm thick, exuding milky latex, leaves broad obovate, elliptic, decurrent, glabrous, entire inflorescence solitary axillaries, cauliflorous and ramiflours on short leafy shoots. Male head is sessile or on short peduncles receptacles, sometimes born on the ultimate twing, Female head are oblong ovoid receptacle, syncarpus, cylindrical (2). Seeds are separated horny endocarpus enclosed by sub-gelatinous exocarpus (1mm thick) oblong ellipsoid in nature. The sweet yellow sheaths around the seeds are about 3-5 mm thick and have a taste similar to that of pineapple, but milder and less juicy (3).

Jackfruit (*Artocarpus heterophyllus* Lam) produces heavier yield than any other tree species, and bear the largest known edible fruit (up to 35 kg). The jackfruit tree has several uses. Flakes of ripe fruits are high in nutritive value; every 100 g of ripe flakes contains 287-323 mg potassium, 30.0-73.2 mg calcium and 11-19 g carbohydrates (4). In Bangladesh, it is commonly referred to as "poor man's food" as it is cheap and plentiful during the season. The nutritious seeds are boiled or roasted and eaten like chestnuts, added to flour for baking, or cooked in dishes. The tree is also known for its durable timber, which ages to an orange or reddish brown color, with anti-termite properties (5). The leaves and fruit waste provide valuable fodder for cattle, pigs and goats. Jackfruit wood chips yield a

dye, which is used to give the famous orange-red color to the robes of Buddhist priests. In addition, many parts of the plant, including the bark, roots, leaves and fruits have medicinal properties (6).

It requires a soil which is well drained but moist, with a ph of 4.3 to 8.0 and with medium soil fertility. The optimum temperature is 19 to 29°C, altitude at approx. 1600 meters above sea level and the annual rain fall between 1000 and 2400 mm (7).

Taxonomical classification (8)

Kingdom	:Plantae-- planta, plantes, plants, vegetal
Subkingdom	:Tracheobionta -- vascular plants
Division	:Magnoliophyta -- angiosperms, flowering plants, phanérogames
Class	:Magnoliopsida -- dicots, dicotylédones, dicotyledons
Subclass	:Hamamelididae
Order	:Urticales
Family	:Moraceae -- mulberries
Genus	:Artocarpus – breadfruit
Species	: <i>Artocarpus heterophyllus</i> Lam.

Parts used:

Apart from whole plant, seeds, fruits, bark, root, leaves and latex are also used.

Synonyms: (9)

Artocarpus brasiliensis Gomez., *Artocarpus heterophylla* Lam., *Artocarpus maxima* Blanco, *Artocarpus philippinensis* Lam., *Polypheuma jaca* Lour., *Soccus arboreus major* Rumph., *Artocarpus integra* (Thunb.), *Artocarpus integrifolia* L.f., *Artocarpus integrifolius* auct., *Artocarpus integer* auct.

Ayurvedic Description

Sanskrit name	:Panasa Synonyms: Atibrhatphala
Properties	: Rasa: Madhura, kasaya (unripe fruit); Guna: Snigdha, guru, picchila; Virya: Sita; Vipaka: Madhura

Artocarpus heterophyllus (Jackfruit): An Overview

Actions :Vatapittahara, kaphavardhaka, balaya, sukraprada, tarpana, durjara,
Seeds- Vrsya, baddhavikara, mutrala
Therapeutic uses :Klibata, durbalya, raktapitta, varna

Growth & Distribution

Artocarpus heterophyllus grows rapidly in early years, up to 1.5 m/yr (5 ft/yr) in height, slowing to about 0.5 m/yr (20 in/yr) as the tree reaches maturity (10). Jackfruit has been cultivated since prehistoric times and has naturalized itself in many parts of the tropics, particularly in Southeast Asia, where it is today an important crop of India, Myanmar, China, Sri Lanka, Malaysia, Indonesia, Thailand and Philippines. It is also grown in parts of Africa, Brazil, Surinam, Caribbean, Florida and Australia. It has been introduced to many Pacific islands since post European contact and is of particular importance in Fiji, where there is a large population of Indian descent.

MORPHOLOGY

Size and form

Jackfruit is a medium size, evergreen tree that typically attains a height of 8–25 m (26–82 ft) and a stem diameter of 30–80 cm (12–32 in). The canopy shape is usually conical or pyramidal in young trees and becomes spreading and domed in older trees. The tree casts a very dense shade. Heavy side branching usually begins near the ground. All parts of the tree exude sticky white latex when injured.

Flowers

This species is monoecious, having male and female inflorescences (or “spikes”) on the same tree. Male and female spikes are borne separately on short, stout stems that sprout from older branches and the trunk. Male spikes are found on younger branches above female spikes. Male spikes are dense, fleshy, cylindrical to club shaped, and up to 10 cm (4 in) in length. Flowers are tiny, pale green when young, turning darker with age. Female flowers are larger, elliptic or rounded, with a tubular calyx. The flowers are reportedly pollinated by insects and wind, with a high percentage of crosspollination.

Leaves

Leaves are dark green, alternate, entire, simple, glossy, leathery, stiff, large (up to 16 cm [6 in] in length), and elliptic to oval in form. Leaves are often deeply lobed when juvenile and on young shoots.

Fruit

Jackfruit has a compound or multiple fruit (syncarp) with a green to yellow brown exterior rind that is composed of hexagonal, bluntly conical carpel apices that cover a thick, rubbery, whitish to yellowish wall. The acid to sweetish (when ripe) banana flavored flesh (aril) surrounds each seed. The heavy fruit is held together by a central fibrous core. Fruits are oblong cylindrical in shape, typically 30–40 cm (12–16 in) in length.

Seeds

Seeds are light brown, rounded, 2–3 cm (0.8–1.2 in) in length by 1–1.5 cm (0.4–0.6 in) in diameter, and enclosed in a thin, whitish membrane. Up to 500 seeds can be found in each

fruit. Seeds are recalcitrant and can be stored up to a month in cool, humid conditions (11).

PHYTOCHEMISTRY

The *Artocarpus heterophyllus* contains various chemical constituents as several flavones colouring matters, morin, dihydromorin, cynomacurin, artocarpin (Fig.2), isoartocarpin, cyloartocarpin, artocarpesin, oxydihydroartocarpesin, artocarpetin, norartocarpetin, cycloartinone and artocarpanone (12). The heart wood on analysis yields moisture 6.7%, glucosides 38.0%, lipids 0.7%, albumin 1.7% and cellulose 59.0 % (13). The plant also contains free sugar (sucrose), fatty acids, ellagic acid and some essential Amino acids like Arginine, Cystine, Histidine, Leucine, Lysine, Methionine, Theonine, Tryptophan etc. (14). Bark from main trunk contains betullic acid and two new flavone pigments, cycloheterophyllin (C₃₀H₃₀O₇) (15). Triterpenic compounds like cycloartenyl acetate, cycloartenone are also reported (16). Heterophyllol (Fig.1) a phenolic compound with novel skeleton was obtained from *Artocarpus heterophyllus* (17). There is only 3.3% tannin in the bark, which is occasionally made into cordage or cloth. The leaves and stem show the presence of sapogenins, cycloartenone, cycloartenol (Fig.4), β-sitosterol (18) and tannins, they show estrogenic activity. A root contains β-sitosterol (Fig.7), ursolic acid, Betulinic acid (Fig.3) and cycloartenone (19).

Jacalin, the major protein from the *Artocarpus heterophyllus* seeds, is a tetrameric two-chain lectin combining a heavy chain of 133 amino acid residues with a light β chain of 20–21 amino acid residues. It is highly specific for the O-glycoside of the disaccharide Thomsen–Friedenreich antigen (Galβ1–3GalNAc), even in its sialylated form. This property has made jacalin suitable for studying various O-linked glycoproteins, particularly human IgA1 (20). Jacalin's uniqueness in being strongly mitogenic for human CD4⁺ T lymphocytes has made it a useful tool for the evaluation of the immune status of patients infected with human immunodeficiency virus HIV-1 (21). Two novel 2', 4', 6'-trioxygenated flavanones, heteroflavanones A and B were isolated from the root bark of *Artocarpus heterophyllus*. Their structures were elucidated as 5-hydroxy-7,2', 4',6'-tetramethoxyflavanone and 8-(γ,γ-dimethylallyl)5-hydroxy-7,2',4',6'-tetramethoxyflavanone (22, 23). Three phenolic compounds were characterized as artocarpesin [(5,7,2',4'-tetrahydroxy-6-(3-methylbut-3-enyl) flavone], norartocarpetin (5,7,2',4'-tetrahydroxyflavone) and oxyresveratrol (Fig.6) (trans-2,4,3',5'-tetrahydroxystilbene) by spectroscopic methods and through comparison with data reported in the literatures (24). The anti-inflammatory effects of these isolated compounds were evaluated by determining their inhibitory effects on the production of proinflammatory mediators in lipopolysaccharide (LPS)-activated RAW 264.7 murine macrophage cells. These three compounds exhibited potent anti-inflammatory activity (25). The composition of carotenoids of *A.heterophyllus* is carotenes β-carotene, α-carotene, β-zeacarotene, α-zeacarotene and β-carotene-5, 6-epoxide and a dicarboxylic carotenoid, crocetin were identified (26).




Fig.1 Heterophyllol




Fig.2 Artocarpin




Fig. 3 Betulinic acid




Fig.4 Cycloartenol




Fig.5 Cycloheterophylline




Fig.6 Oxyresveratrol




Fig. 7 β -Sitosterol

Traditional uses

The leaves are useful in fever, boils, wounds and skin diseases. The young fruits are acrid, astringent, and carminative. The ripe fruits are sweet, cooling, laxative, aphrodisiac and also used as a brain tonic. The seeds are, diuretic, and constipating. The wood is nervine, antidiabetic, sedative and is useful in convulsions (27). The latex is useful in dysopia, ophthalmic disorders and pharyngitis and also used as antibacterial agent (28). The ash of Jackfruit leaves is used in case of ulcers. The dried latex yields artostenone, convertible to artosterone, and a compound with marked androgenic action. Mixed with vinegar, the latex promotes healing of abscesses, snakebite and glandular swellings (29). The root is a remedy for skin diseases and asthma. An extract of the root is taken in cases of fever and diarrhoea. The bark is made into poultices. Heated leaves are placed on wounds. The wood has a sedative property and its pith is said to be abortifacient. Latex is used as an anti-inflammatory agent (30).

PHARMACOLOGY

Although a lot of pharmacological investigations have been carried out based on the constituents present in it but a lot more can still be explored and utilized in a therapeutic manner. A summary of the findings of some of these studies is presented below.

Anti-inflammatory Effect

Artocarpus heterophyllus Lam is a large evergreen tree cultivated throughout Southeast Asia for its fruits. Its leaves and roots have been used for medicinal purposes. The aim of this work was to study the *in vitro* anti-inflammatory effects of phenolic compounds isolated from the ethyl acetate extracts of the fruits of *Artocarpus heterophyllus*. Three phenolic compounds were characterized as artocarpesin [5,7,2',4'-tetrahydroxy-6-(3-methylbut-3-enyl) flavone] (1), norartocarpetin (5,7,2',4'-tetrahydroxyflavone) (2), and oxyresveratrol [trans-2,4,3',5'-tetrahydroxystilbene] (3) by spectroscopic methods and through comparison with data reported in the literatures. The anti-inflammatory effects of the isolated compounds (1- 3) were evaluated by determining their inhibitory effects on the production of proinflammatory mediators in lipopolysaccharide (LPS)-activated RAW 264.7 murine macrophage cells. These three compounds exhibited potent anti-inflammatory activity. The results indicated that artocarpesin (1) suppressed the LPS-induced production of nitric oxide (NO) and prostaglandin E 2 (PGE 2) through the down-regulation of inducible nitric oxide synthase (iNOS) and cyclooxygenase 2 (COX-2) protein expressions. Thus, artocarpesin (1) may provide a potential therapeutic approach for inflammation-associated disorders (31).

Artocarpus heterophyllus (Jackfruit): An Overview**Antioxidant Effect**

The antioxidant properties of prenylflavones, isolated from *Artocarpus heterophyllus* Lam., were evaluated in this study. Among them, artocarpine, artocarpetin, artocarpetin A, and cycloheterophyllin diacetate and peracetate had no effect on iron-induced lipid peroxidation in rat brain homogenate. They also did not scavenge the stable free radical 1, 1-diphenyl-2-picrylhydrazyl. In contrast, cycloheterophyllin (Fig. 5) and artonins A and B inhibited iron-induced lipid peroxidation in rat brain homogenate and scavenged 1, 1-diphenyl-2-picrylhydrazyl. They also scavenged peroxy radicals and hydroxyl radicals that were generated by 2, 2'-azobis (2-amidinopropane) dihydrochloride and the Fe³⁺-ascorbate-EDTA-H₂O₂ system, respectively. However, they did not inhibit xanthine oxidase activity or scavenge superoxide anion, hydrogen peroxide, carbon radical, or peroxy radicals derived from 2,2'-azobis(2,4-dimethylvaleronitrile) in hexane. Moreover, cycloheterophyllin and artonins A and B inhibited copper-catalyzed oxidation of human low-density lipoprotein, as measured by fluorescence intensity, thiobarbituric acid-reactive substance and conjugated-diene formations and electrophoretic mobility. It is concluded that cycloheterophyllin and artonins A and B serve as powerful antioxidants against lipid peroxidation when biomembranes are exposed to oxygen radicals (32).

Antifungal Effect

Two novel chitin-binding lectins from seeds of *Artocarpus* genus were described, one from jackfruit and one from breadfruit. They were purified from saline crude extract of seeds using affinity chromatography on chitin column, size-exclusion chromatography and reverse-phase chromatography on the C-18 column. Both are 14 kDa proteins, made up of 3 chains linked by disulfide bonds. The partial amino acid sequences of the two lectins showed they are homologous to each other but not to other plant chitin-binding proteins. Thus, they cannot be classified in any known plant chitin-binding protein family, particularly because of their inter-chain covalent bonds. Their circular dichroism spectra and deconvolution showed a secondary structure content of beta-sheet and unordered elements. The lectins were thermally stable until 80 degrees C and structural changes were observed below pH 6. Both lectins inhibited the growth of *Fusarium moniliforme* and *Saccharomyces cerevisiae*, and presented hemagglutination activity against human and rabbit erythrocytes. These lectins were denoted jackin (from jackfruit) and frutackin (from breadfruit) (33).

Sexual behaviour

According to medicinal plants text of Sri Lanka, roasted seeds of *Artocarpus heterophyllus* Lam. (Family: Moraceae) has aphrodisiac activity. However, some reproductively active young men in rural areas of Sri Lanka claim that consumption of these seeds few hours prior to coitus disrupts sexual function. Because of these two conflicting claims, it was thought useful to scientifically investigate the effects of *A. heterophyllus* seeds on male sexual function and fertility. This was done using a seed suspension in 1% methylcellulose (SS) in rats. In a sexual behaviour study using receptive female rats,

an oral administration of 500 mg/kg dose of SS markedly inhibited libido, sexual arousal, sexual vigour and sexual performance within 2 hr. Further, the treatment induced a mild erectile dysfunction. These antimasculine effects on sexual function were not evident 6 hr post treatment indicating rapid onset and offset of action. Further, these actions on the sexual behaviour was not due to general toxicity, liver toxicity, stress or reduction in blood testosterone level but due to marked sedative activity. In a mating study, SS failed to alter ejaculating competence and fertility. These results suggest that *A. heterophyllus* seeds do not have aphrodisiac action, at least, in rats (34).

Immunomodulatory effect

Jacalin, the major protein from the jackfruit (*Artocarpus heterophyllus*) seeds, is a tetrameric two-chain lectin (molecular mass 65 kDa) combining a heavy chain of 133 amino acid residues with a light β chain of 20–21 amino acid residues. It is highly specific for the O-glycoside of the disaccharide Thomsen–Friedenreich antigen (Gal β 1–3GalNAc), even in its sialylated form. This property has made jacalin suitable for studying various O-linked glycoproteins, particularly human IgA1. Jacalin's uniqueness in being strongly mitogenic for human CD4⁺ T lymphocytes has made it a useful tool for the evaluation of the immune status of patients infected with human immunodeficiency virus (HIV)-1. The abundance of source material for the production of jacalin, its ease of purification, yield and stability has made it an attractive cost-effective lectin. It has found applications in diverse areas such as the isolation of human plasma glycoproteins (IgA1, C1-inhibitor, hemopexin, 2-HSG), the investigation of IgA-nephropathy, the analysis of O-linked glycoproteins and the detection of tumours (35).

Antidiabetic Effect

Investigations were carried out to evaluate the effects of hot-water extract of *Artocarpus heterophyllus* leaves on the glucose tolerance of normal human subjects and maturity-onset diabetic patients. The extracts of *Artocarpus heterophyllus* significantly improved glucose tolerance in the normal subjects and the diabetic patients when investigated at oral doses equivalent to 20 g/kg of starting material (36).

Antibacterial Effect

The crude methanolic extracts of the stem and root, barks, stem and root heart-wood, leaves, fruits and seeds of *Artocarpus heterophyllus* and their subsequent partitioning with petrol, dichloromethane, ethyl acetate and butanol gave fractions that exhibited a broad spectrum of antibacterial activity. The butanol fractions of the root bark and fruits were found to be the most active (37).

Anthelmintic Effect

The shoots revealed nematicidal activity against various nematodes viz., *Rotylenchulus reniformis*, *Tylenchorhynchus brassicae*, *Tylenchus filiformis* and *Meloidogyne incognita* (38).

Inhibition of melanin biosynthesis

A series of prenylated, flavones based polyphenols, compounds 1-8, and were isolated from the wood of *Artocarpus heterophyllus*. These compounds, which have previously been shown not to inhibit tyrosinase activity, were

found to be active inhibitors of the in vivo melanin biosynthesis in B16 melanoma cells, with little or no cytotoxicity. To clarify the structural requirement for inhibition, some structure-activity relationships were studied, in comparison with related compounds lacking prenyl side chains. Our experiments indicate that both prenyl and OH groups, as well as the type of substitution pattern, are crucial for the inhibition of melanin production in B₁₆ melanoma cells (39).

Allergy to jackfruit

Jackfruit allergy has been reported just once. It is unknown whether this food allergy is caused by direct sensitization or cross-sensitization to pollen allergens. Jackfruit allergy can be added to the list of birch pollen-related food allergies. Increased consumption of this fruit will result in a rise in allergic reactions (40).

Some chemical constituents isolated from *A. heterophyllum*

Six prenylflavonoids, including two new compounds, have been isolated from the root, bark of *Artocarpus heterophyllum*. The new prenylflavones have been characterized as 8-(γ,γ -dimethylallyl)-5,4'-dihydroxy-7,2'-dimethoxyflavone and 3,3'-di-(γ,γ -dimethylallyl)-5,7,2',5'-tetrahydroxy-4'-methoxyflavone, respectively (41).

A new natural Diels-Alder-type adduct, artonin X, along with two known Diels-Alder type adducts, were isolated from the bark of *Artocarpus heterophyllum* (42).

In an investigation emission and Fourier transform infrared spectra of the jackfruit seed, in powdered form, were recorded. Analysis of the emission spectrum confirms the presence of two hitherto undetected elements, manganese and magnesium. The Fourier transform infrared spectrum reveals the presence of some specific functional groups, attributed to the different bands present in the spectrum (43).

PESTS AND DISEASES

In southwestern and southern Asia, boring insects seem to be the major pests of jackfruit. These include *Indarbela tetraonis*, *Batocera rufomaculata*, *Margaronia caecalis*, and *Ochyromera artocarpio*. In India the main insect pests are the shoot boring caterpillar (*Diaphania caecalis*), mealybugs (*Nipaecoccus viridis*, *Pseudococcus corymbatus*, and *Ferrisia virgata*), spittle bug (*Cosmoscarta relata*) and jack scale (*Ceroplastes rubina*). In southern China, the fruit stem is susceptible to damage from the larvae of the longicorn beetles *Aprona germari*, *Pterolophia discalis*, *Xenolea tomentosa asiatica*, and *Olenecamptus bilobus*. The caterpillars of leaf webbers (*Perina nuda* and *Diaphania bivitalis*), aphids (*Greenidea artocarpis* and *Toxoptera aurantii*), and thrips (*Pseudodendrothrips dwivarna*) are minor problems. Important diseases of jackfruit are pink disease (*Pelliculana* [syn. *Corticium*] *salmonicolor*); stem, fruit, and male inflorescence rot caused by *Rhizopus artocarpis*; and leafspot due to *Phomopsis artocarpina*, *Colletotrichum lagenarium*, *Septoria artocarpis*, and other fungi. Gray blight (*Pestalotia elasticola*), charcoal rot (*Ustilana zonata*), collar rot (*Rosellinia arcuata*), and rust (*Uredo artocarpis*) occur on jackfruit in some regions.

In a recent study it was found that wood boring insects include *Elaphidion mucronatum*, *Nyssodrycina baldemani*, and

Leptostylopsis terraeicolor various scales and mealybugs may attack stems and fruit. Diseases include *Rhizopus* fruit rot (*Rhizopus artocarpis*), gray mold (*Botrytis cinerea*), root rot (*Pythiumsplendens*, *Phytophthora* sp., *Fusarium* sp., and *Rhizoctonia* sp.), and leaf spotting by fungi (*Gloeosporium* sp. and *Phyllotacticia artocarpis*) also affect the plant (44).

CONCLUSION

Considering the easy availability of *Artocarpus heterophyllum* in our country and that almost all the parts of the plant, including wood & latex possess curative properties, it seems that still there is a scope for scientific studies to fully exploit its medicinal properties to support the traditional claims as well as ,exploring some new and promising 'leads'. This review is an effort to compile all major information on its phytochemical as well as pharmacological profile published till now.

REFERENCES

1. A.M. Rahman, N. Nahar, A.J. Mian and M. Mosihuzzaman. Variation of carbohydrate composition of two forms of fruit from jack tree (*Artocarpus heterophyllum* L.) with maturity and climatic conditions. *Food Chem.* **65**: 91-97 (1999).
2. P. Rowe-Dutton. *Artocarpus heterophyllum*- jackfruit. In: The propagation of tropical fruit trees (Garner RJ and Chaudhri SA, eds.). FAO, Rome (Italy); Commonwealth Bureau of Horticulture and Plantation Crops, Maidstone, 269-290 (1985).
3. The Wealth of India, *A dictionary of Indian raw materials and industrial products*, (publication and information directorate CSIR, New Delhi, 1985), 445-453.
4. C. E. Elevitch, and H. I. Manner. *Artocarpus heterophyllum* (jackfruit). Species Profiles for Pacific Island Agroforestry (2006).
5. E. T. Arung, K. Shimizu and R. Kondo. Inhibitory effect of artocarpinone from *Artocarpus heterophyllum* on melanin biosynthesis. *Biol. Pharm. Bull.* **29**: 1966-1969 (2006).
6. S. Shyamamma, S.B.C. Chandra, M. Hegde and P. Naryanswamy. Evaluation of genetic diversity in jackfruit (*Artocarpus heterophyllum* Lam.) based on amplified fragment length polymorphism markers. *Genet. Mol. Res.* **7**(3): 645-656 (2008).
7. Roy, S.K., P.K. Roy and R.G. Brumfield, *In vitro* propagation and establishment of a new cultivar of jackfruit (*Artocarpus heterophyllum* lam.) bearing fruits twice yearly. *Acta Hort.*, **429**: 497-502 (1996).
8. <http://plants.usda.gov/java/ClassificationServlet?source=display&classid=ARHE2>
9. <http://www.agroforestry.net/tti/A.heterophyllum-jackfruit.pdf>
10. M. Kamaluddin, M. Ali and M.K. Bhuiyan. Effect of auxin on rooting of cuttings and growth of seedlings of jackfruit (*Artocarpus heterophyllum* lam.). *Chittagong Univ. Stud. Sci.* **20**(1): 71-75(1997).
11. www.traditionaltree.org
12. A.V. Rama Rao, Mala Varadan and Venkataraman. Colouring matter of the *A. heterophyllum*. *Indian J. Chem.* **11**: 298-299 (1973).
13. Perkin and Cope. The constituents of *Artocarpus integrifolia*. *J. Chem. Soc.* **67**: 937-944 (1895).
14. G. Pavanadasivam, M. Uvais and S. Sultanbawa. Cycloartenyl acetate, cycloartenol and cycloartenone in the bark of *Artocarpus species*. *Phytochemistry*, **12**(11): 2725-2726 (1973).
15. F.A. Chawdhary, M.A. Raman. Distribution of free sugars and Fatty acids in Jackfruit. *Food chemistry.* **60**(1): 25-28 (1997).
16. B.R. Barik, T. Bhaumik, A.K. and A.B. Kundu. Triterpenoids of *Artocarpus heterophyllum*, *J. Indian Chemical Soc.* **74**: 163-164 (1997).
17. Chun-Nan Lin and Chai-Ming Lu. Heterophyllol, a phenolic compound with novel skeleton from *Artocarpus heterophyllum*, *Tetrahedron letters.* **34**(17): 8249-8250 (1993).
18. M.C. Nath, 1937b. *Z. Physiology Chem.* **247**, 9. Quoted in Plant, and Chaturvedi, K. *Phytochemistry* **28**: 2197-2199 (1989).
19. R. Dayal and T.R. Seshadri. Colourless compounds of the roots of *Artocarpus heterophyllum*. Isolation of new compound artoflavone. *Indian J Chem.* **12**: 895-896 (1974).

Artocarpus heterophyllus (Jackfruit): An Overview

20. G. Suresh Kumar, P.S. Appuktan and D.K. Basu. D-Galactose – specific lectin from jack fruit seed. *J. Biosci.* **4**: 257-261 (1982).
21. G. Pereira-da-Silva, AN. Moreno, F. Marques, C. Oliver, MC. Jamur, A. Panunto-Castelo, MC. Roque-Barreira, Neutrophil activation induced by the lectin KM+ involves binding to CXCR2. *Biochim. Biophys. Acta.* **1**: 86-94 (2006).
22. Chai-Ming Lu and Chun-Nan Lin. Two 2', 4', 6'-trioxygenated flavanones from *Artocarpus heterophyllus*. *Natural Products Research Center* **33(4)**: 909-911 (1993).
23. Chun-Nan Lin, Chai-Ming Lu and Pao-Lin Huang. Flavonoids from *Artocarpus heterophyllus*. *Phytochemistry*. **39(6)**: 1447-1451 (1995).
24. K. Venkataraman. Wood of phenolics in the chemotaxonomy of the Moraceae. *Phytochemistry* **11(5)**: 1571-1586 (2001).
25. P.K. Jha, R.S. Choudhary, and S.K. Choudhary. Studies of medicinal plants of Palamau (Bihar)-IInd Part. *Biojournal* **9**: 21-38 (1997).
26. U.G. Chandrika, E.R. Jansz and N.D. Warnasuriya. Analysis of carotenoids in ripe jackfruit (*Artocarpus heterophyllus*) kernel and study of their bioconversion in rats. *Journal of the Science of Food and Agriculture* **85(2)**: 186 -190 (2004).
27. P. P. Hemborn. Contact therapy practiced by Mundas Chotanagar (Bihar). *Ethnobotany*. **8**: 36-39 (1996).
28. M. Sato and S. Fujiwara. Flavones with Antibacterial activity against carcinogenic bacteria. *J. Ethnopharmacol.* **54(2-3)**: 171-176 (1996).
29. V. M. Vaidya Gogte, *Ayurvedic Pharmacology and therapeutic use of medicinal plants*, (Swami Prakashananda Ayurvedic Research center, Mumbai, 2000) 656-657.
30. A. K. Gupta and N. Tandon, *Review on Indian Medicinal Plants*, (Indian Council of Medical Research, New Delhi, 2004) 182-200.
31. S.C. Fang, C.L. Hsu, G.C. Yen. Anti-inflammatory effects of phenolic compounds isolated from the fruits of *Artocarpus heterophyllus*. *J Agric Food Chem.* **56(12)**: 4463-4468 (2008).
32. F.N. Ko, Z.J. Cheng, C.N. Lin and C.M. Teng. Scavenger and antioxidant properties of prenylflavones isolated from *Artocarpus heterophyllus*. *Free Radic Biol Med.* **25(2)**: 160-168 (1998).
33. M.B. Trindade, J.L. Lopes, A. Soares Costa, A.C. Monteiro Moreira, R.A. Moreira, M.L. Oliva and L.M. Beltramini. Structural characterization of novel chitin-binding lectins from the genus *Artocarpus* and their antifungal activity. *Biochim Biophys Acta.* **1764(1)**: 146-52 (2006).
34. W.D. Ratnasooriya, J.R. Jayakody. *Artocarpus heterophyllus* seeds inhibit sexual competence but not fertility of male rats. *Indian J Exp Biol.* **40(3)**: 304-308 (2002).
35. S. Kabir. Jacalin, a jackfruit (*Artocarpus heterophyllus*) seed-derived lectin of versatile applications in Immunobiological research. *Journal of Immunological Method* **212(2)**: 193-211 (1998).
36. M. R. Fernando, S. M. D. Nalinie Wickramasinghe, M. I. Thabrew, P. L. Ariyananda and E. H. Karunanayake. Effect of *Artocarpus heterophyllus* and *Asteracanthus longifolia* on glucose tolerance in normal human subjects and in maturity-onset diabetic patients. *J. Ethnopharmacol.* **31(3)**: 277-282 (1991).
37. M. R. Khan, Omoloso and M. Kihara. Antibacterial activity of *Artocarpus heterophyllus*. *Fitoterapia* **74(5)**: 501-505 (2003).
38. W. Sharma and P.C. Trivedi. Nematocidal and nematostostatic response of aqueous extract of certain plants of semi arid niche. *Curr. Nematol.* **6**: 45-53 (1995).
39. E.T. Arung, K. Shimizu and R. Kondo. Structure-activity relationship of prenyl-substituted polyphenols from *Artocarpus heterophyllus* as inhibitors of melanin biosynthesis in cultured melanoma cells. *Chem Biodivers* **4(9)**: 2166-71 (2007).
40. S.T. Bolhaar, R. Ree, C.A. Buijnzeel-Koomen, A.C. Knulst and L. Zuidmeer. Allergy to jackfruit, a novel example of Bet v 1-related food allergy. *Allergy.* **59(11)**: 1187-92 (2004).
41. M. Chung, C. Lu, P. Huang, and C. Lin. Prenylflavonoids of *Artocarpus heterophyllus*. *Phytochemistry.* **40(4)**: 1279-1282 (1995).
42. K. Shinomiya, M. Aida, Y. Hano and T. Nomura. A diels-alder-type adducts from *Artocarpus heterophyllus*. *Phytochemistry.* **40(4)**: 1317-1319 (1995).
43. A.G. Barua, B.R. Boruah. Minerals and functional groups present in the jackfruit seed: a spectroscopic investigation. *J. Food Sci. Nutr.* **55(6)**: 479-83 (2004).
44. J. Morton, F. Julia, Morton and Miami. *In Fruits of warm climates.* 58-64 (1987).